# This Page Is Inserted by IFW Operations and is not a part of the Official Record

### BEST AVAILABLE IMAGES

Defective images within this document are accurate representation of The original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT.
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

### IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE BEFORE THE BOARD OF PATENT APPEALS AND INTERPERENCES

JEAN-YVES CHENARD et al.

Serial No.: 07/870,759 ) Group Art Unit: 1511
Filed: April 20, 1992 ) Examiner: V. Hoke
For: IMPROVEMENT IN THE )
STABILIZATION OF VINYL )
HALIDE POLYMERS

#### CLAIMS ON APPPAL

- 176. A method of stabilizing a vinyl halide resin comprising the steps of:
- (a) adding to the vinyl halide resin a mono- or diorganotin compound wherein at least one atom bonded to tin is a halogen; and
- (b) adding to the vinyl halide resin a mercapto alkanol ester of a monocarboxylic acid;

the mono-or diorganotin compound and the mercapto alkanol ester of a monocarboxylic acid being present in an amount effective to stabilize the vinyl halide resin.

177. The method of claim 176, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

#### RCOOR'SH

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 178. The method of claim 177, wherein R contains 6 to 38 carbon atoms.
- 179. The method of claim 177, wherein R contains 8 to 18 carbon atoms.

FOREST TEST

FOREST HENDEADN

FOREST CARRETT

SOUNCE

FOR NOTE TO FOREST

FOR NOTE TO FOREST

FOREST ADD FOR

180. The method of claim 179, wherein R' contains 2 to 6 carbon atoms.

- 181. The method of claim 175, wherein the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.
- 182. The method of claim 176, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate, and mercapto ethyl myristate.
- 183. The method of claim 176, wherein the vinyl halide resin is polyvinyl chloride.
- 193. The composition of claim 237, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 194. The composition of claim 237, wherein R contains 6 to 38 carbon atoms.
- 195. The composition of claim 237, wherein R contains 8 to 18 carbon atoms.
- 196. The composition of claim 237, wherein R contains 2 to 6 carbon atoms.
- 197. The composition of claim 237, the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.

FINALIAN HENDERSON FRANCISCH CARRETT 6 DINNER 1000 1 STREET N. L. TEST NITTON DI 20005 1 2000

JEAN-IVES CHENARD et al. Serial No. 07/870,759

- 198. The composition of claim 237, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.
- 200. The vinyl halide resin composition of claim 239, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

#### RCOOR'SH

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 201. The vinyl halide resin composition of claim 239, wherein R contains 6 to 38 carbon atoms.
- 202. The vinyl halide resin composition of claim 239, wherein R contains 8 to 18 carbon atoms.
- 203. The vinyl halide resin composition of claim 239, wherein R contains 2 to 6 carbon atoms.
- 204. The vinyl halide resin composition of claim 239, the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.
- 205. The vinyl halide resin composition of claim 239, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.

JEAN-YVES CHENARD et al. Serial No. 07/870,759

- 206. The vinyl halide resin composition of claim 239, wherein the mercapto alkanol ester of a monocarboxylic acid is present in the range of 0.1 % wt. to 5 % wt. of the vinyl halide resin.
- 207. The vinyl halide resin composition of claim 239, wherein the vinyl halide resin is polyvinyl chloride.
- 209. The composition of claim 241, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

#### RCOOR'SH

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $C_2$  to  $C_{1,2}$  alkylene.

- 210. The composition of claim 241, wherein R contains 6 to 38 carbon atoms.
- 211. The composition of claim 241, wherein R contains 2 to  $\epsilon$  carbon atoms.
- 212. The composition of claim 241, the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.
- 213. The composition of claim 241, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.
- 214. The composition of claim 241, further comprising a vinyl halide resin.

FINEDAY HENDERSON
FARACCII CARRETT

É DINNER

100: STREET N W
PARACCII STREET N W

- 215. The composition of claim 241, wherein the product produced by mixing the organotin compound and the mercapto alkanol ester of a monocarboxylic acid are present in an amount effective to stabilize the vinyl halide resin against heat or light.
- 216. The composition of claim 241, wherein the mercapto alkanol ester of a monocarboxylic acid is present in the range of 0.1 % wt. to 5 % wt. of the vinyl halide resin.
- 217. The composition of claim 241, wherein the vinyl halide resin is polyvinyl chloride.
- 219. The method of claim 243, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 220. The method of claim 243, wherein R contains 6 to 38 carbon atoms.
- 221. The method of claim 243, wherein R contains 8 to 18 carbon atoms.
- 222. The method of claim 243, wherein R contains 2 to 6 carbon atoms.
- 223. The method of claim 243, the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.

FINNEDAN, HENDEADON
FARABOR GARRETT

6 DINNER
1000 SEPRET N. 4
7454-100 C 20005
1-202-403-4000

JEA VES CHENARD et al. Serial No. 07/870,759

- 224. The method of claim 243, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.
- 225. The method of claim 243, wherein the vinyl halide resin is polyvinyl chloride.
- 227. The method of claim 245, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

#### RCOOR'SH

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 228. The method of claim 245, wherein R contains 6 to 38 carbon atoms.
- 229. The method of claim 245, wherein R contains 8 to 18 carbon atoms.
- 230. The method of claim 245, wherein R contains 2 to 6 carbon atoms.
- 231. The method of claim 245, the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.
- 232. The method of claim 245, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.
- 233. The method of claim 245, wherein the vinyl halide resin is polyvinyl chloride.

FORMED N. HENDERSON FARAGOS, CARRETT 5 DINNER 100 1 5 76 CT & 2 TOSANDOS DO ECOS

- 237. A composition comprising:
- (a) a mono- or dialkyltin bis(isooctyl mercapto acetate) compound wherein the alkyl is a  $C_1$  to  $C_8$  alkyl; and
  - (b) a mercapto alkanol ester of a monocarboxylic acid.
- 238. A composition according to claim 237, wherein said mono- or dialkyltin bis(isooctyl mercapto acetate) compound is selected from the group consisting of di-n-octyltin bis-(isooctyl mercapto acetate); di-n-butyltin bis-(isooctyl mercapto acetate); a mixture of an anhydride of thiobutyl stannic acid with di-n-butyltin bis-(isodecyl mercapto acetate); and a condensation polymer of butyl stannic acid and butyl thiostannic acid.
  - 239. A vinyl halide resin composition comprising:
  - (a) a vinyl halide resin;
- (b) a mono- or dialkyltin bis(isooctyl mercapto acetate) compound wherein the alkyl is a  $C_1$  to  $C_8$  alkyl; and
  - (c) a mercapto alkanol ester of a monocarboxylic acid.
- 240. A composition according to claim 239, wherein said mono- or dialkyltin bis(isooctyl mercapto acetate) compound is selected from the group consisting of di-n-octyltin bis-(isooctyl mercapto acetate); di-n-butyltin bis-(isooctyl mercapto acetate); a mixture of an anhydride of thiobutyl stannic acid with di-n-butyltin bis-(isodecyl mercapto acetate); and a condensation polymer of butyl stannic acid and butyl thiostannic acid.
  - 241. A composition comprising a product produced by mixing:
- (a) a mono- or dialkyltin bis(isooctyl mercapto acetate) compound wherein the alkyl is a  $C_1$  to  $C_8$  alkyl; and
  - (b) a mercapto alkanol ester of a monocarboxylic acid.

PONICAN MINDERSON
FARADOR DARATTO
S DINNER
DD 1 STACTOR A
TASHADOR DE 20005
A 202 A 20 A 202

- 242. A composition according to claim 241, wherein said mono- or dialkyltin bis(isooctyl mercapto acetate) compound is selected from the group consisting of di-n-octyltin bis-(isooctyl mercapto acetate); di-n-butyltin bis-(isooctyl mercapto acetate); a mixture of an anhydride of thiobutyl stannic acid with di-n-butyltin bis-(isodecyl mercapto acetate); and a condensation polymer of butyl stannic acid and butyl thiostannic acid.
- 243. A method of stabilizing a vinyl halide resin comprising the steps of:
- (a) adding to the vinyl halide resin a mono- or dialkyltin bis(isooctyl mercapto acetate) compound wherein the alkyl is a  ${\bf C}_1$  to  ${\bf C}_8$  alkyl; and
- (b) adding to the vinyl halide resin a mercapto alkanol ester of a nonocarboxylic acid.
- 244. A method according to claim 243, wherein said mono- or dialkyltin bis(isooctyl mercapto acetate) compound is selected from the group consisting of di-n-octyltin bis-(isooctyl mercapto acetate); di-n-butyltin bis-(isooctyl mercapto acetate); a mixture of an anhydride of thiobutyl stannic acid with di-n-butyltin bis-(isodecyl mercapto acetate); and a condensation polymer of butyl stannic acid and butyl thiostannic acid.
- 245. A method of stabilizing a vinyl halide resin comprising the step of adding to the vinyl halide resin in an amount effective to stabilize the vinyl halide resin a product produced by mixing:

FINNEAN, HINDEASON
FARAGOR CARRETT
É DINNER
1900 I STREET N =
74871N0TON DI 20003
1 2014 408 4000

- (a) a mono- or dialkyltin bis(isooctyl mercapto acetate) compound wherein the alkyl is a  $\rm C_1$  to  $\rm C_8$  alkyl; and
  - (b) a mercapto alkanol ester of a monocarboxylic acid.
- 246. A method according to claim 245, wherein said mono- or dialkyltin bis(isooctyl mercapto acetate) compound is selected from the group consisting of di-n-octyltin bis-(isooctyl mercapto acetate); di-n-butyltin bis-(isooctyl mercapto acetate); a mixture of an anhydride of thiobutyl stannic acid with di-n-butyltin bis-(isodecyl mercapto acetate); and a condensation polymer of butyl stannic acid and butyl thiostannic acid
- in escents effective to stabilize vinyl halide resins:
- (i) a mono- or diorganotin compound wherein at least one atom bonded to tin is sulfur; and
  - (ii) a mercapto alkanol ester of a monocarboxylic acid.
- 248. The composition of claim 247, wherein in the mono- or di- organotin compound there is bonded to tin at least one alkyl mercapto ester group.
- 249. The composition of claim 247, wherein at least one sulfur bonded to tin is the residue of a mercaptan.
- 250. The composition of claim 247, wherein at least one sulfur bonded to tin is the residue of a mercapto acid.
- 251. The composition of claim 247, wherein at least one sulfur bonded to tin is the residue of a mercapto alcohol.
- 252. The composition of claim 247, wherein at least one sulfur bonded to tin is the residue of a mercapto acid ester.
- 253. The composition of claim 247, wherein at least one sulfur bonded to tin is the residue of a mercapto alcohol ester.

254. The composition of claim 247, wherein the mono- or diorganotin compound is selected from the group consisting of di-noctyltin bis-(isooctyl mercapto acetate); di-n-butyltin bis(isooctyl mercapto acetate); a mixture of an anhydride of
thiobutyl stannic acid with di-n-butyltin bis-(isodecyl mercapto
acetate); and a condensation polymer of butyl stannic acid and
butyl thiostannic acid.

255. The composition of claim 247, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

#### RCOOR'SH

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

256. The composition of claim 255, wherein R contains 6 to 38 carbon atoms.

257. The composition of claim 256, wherein R contains 8 to 18 carbon atoms.

258. The composition of claim 255, wherein R' contains 2 to 6 carbon atoms.

259. The composition of claim 247, wherein the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.

260. The composition of claim 247, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.

FINITES FINITES
FINITES HENDERSON
FARAGOR CARACTT

1 DINNER

110 1 STATE N 4
THE NOTE OF THE NOTES

- 261. The composition of claim 247, further comprising a vinyl halide resin.
- 262. The composition of claim 261, wherein the product produced by mixing the organotin compound and the mercapto alkanol ester of a monocarboxylic acid are present in an amount effective to stabilize the vinyl halide resin against heat or light.
- 263. The composition of claim 261, wherein the mercapto alkanol ester of a monocarboxylic acid is present in the range of 0.1 % wt. to 5 % wt. of the vinyl halide resin.
- 264. The composition of claim 261, wherein the vinyl halide resin is polyvinyl chloride.
- 265. A method of stabilizing a vinyl halide resin comprising adding to the vinyl halide resin in an amount effective to stabilize the vinyl halide resin a product produced by mixing:
- (i) a mono- or diorganotin compound wherein at least one atom bonded to tin is sulfur: and
  - (ii) a mercapto alkanol ester of a monocarboxylic acid.
- 266. The method of claim 265, wherein in the mono- or diorganotin compound there is bonded to tin at least one alkyl mercapto ester group.
- 267. The method of claim 265, wherein at least one sulfur bonded to tin is the residue of a mercaptan.
- 268. The method of claim 265, wherein at least one sulfur bonded to tin is the residue of a mercapto acid.
- 269. The method of claim 265, wherein at least one sulfur bonded to tin is the residue of a mercapto alcohol.
- 270. The method of claim 265, wherein at least one sulfur bonded to tin is the residue of a mercapto acid ester.

FINNEDAN HENDERSON
FARADOR CARRETT
6 DINNER
1000 : STREET & B

- 271. The method of claim 265, wherein at least one sulfur bonded to tin is the residue of a mercapto alcohol ester.
- 272. The method of claim 265, wherein the mono- or diorganotin compound is selected from the group consisting of di-noctyltin bis-(isooctyl mercapto acetate); di-n-butyltin bis(isooctyl mercapto acetate); a mixture of an anhydride of
  thiobutyl stannic acid with di-n-butyltin bis-(isodecyl mercapto
  acetate); and a condensation polymer of butyl stannic acid and
  butyl thiostannic acid.
- 273. The method of claim 265, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 274. The method of claim 273, wherein R contains 6 to 38 carbon atoms.
- 275. The method of claim 274, wherein R contains 8 to 18 carbon atoms.
- 276. The method of claim 273, wherein R' contains 2 to 6 carbon atoms.
- 277. The method of claim 265, wherein the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.

FINNEGAN, HENDERSON FARASOW, CARRETT 6 DINNER 1900 I STEET N # TESTINGTON DI 20005 1 202 405 4000 JEAN-YVES CHENARD et al. Serial No. 07/870,759

- 278. The method of claim 265, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.
- 279. The method of claim 265, wherein the vinyl halide resin is polyvinyl chloride.
- 280. In a composition for stabilizing a vinyl halide resin containing a mono- or diorganotin compound wherein at least one atom bonded to tin is sulfur, the improvement comprising the addition of a mercapto alkanol ester of a monocarboxylic acid.
- 281. The composition of claim 280, wherein in the mono- or di- organotin compound there is bonded to tin at least one alkyl mercapto ester group.
- 282. The composition of claim 280, wherein at least one sulfur bonded to tin is the residue of a mercaptan.
- 283. The composition of claim 280, wherein at least one sulfur bonded to tin is the residue of a mercapto acid.
- 284. The composition of claim 280, wherein at least one sulfur bonded to tin is the residue of a mercapto alcohol.
- 285. The composition of claim 280, wherein at least one sulfur bonded to tin is the residue of a mercapto acid ester.
- 286. The composition of claim 280, wherein at least one sulfur bonded to tin is the residue of a mercapto alcohol ester.
- 287. The composition of claim 280, wherein the mono- or diorganotin compound is selected from the group consisting of di-noctyltin bis-(isooctyl mercapto acetate); di-n-butyltin bis(isooctyl mercapto acetate); a mixture of an anhydride of
  thiobutyl stannic acid with di-n-butyltin bis-(isodecyl mercapto

JEAN VES CHENARD et al. Serial No. 07/870,759

acetate); and a condensation polymer of butyl stannic acid and butyl thiostannic acid.

288. The composition of claim 280, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

#### RCOOR'SH

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 289. The composition of claim 288, wherein R contains 6 to 38 carbon atoms.
- 290. The composition of claim 289, wherein R contains 8 to 18 carbon atoms.
- 291. The composition of claim 288, wherein  $\mathbb{R}^2$  contains 2 to 6 carbon atoms.
- 292. The composition of claim 280, wherein the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.
- 293. The composition of claim 280, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.
- 294. The composition of claim 280, wherein the mercapto alkanol ester of a monocarboxylic acid is present in the range of 0.1 % wt. to 5 % wt. of the vinyl halide resin to be stabilized.
- 295. The composition of claim 280, wherein the vinyl halide resin is polyvinyl chloride.

FINNELAN, HENDERSON
FARASON: CARRETT
6 DINNER
1000: STREET N. 6
THANDON DO 20005
1 202 404 4000

JEAN-LES CHENARD et al. Serial No. 07/870,759

- 296. A composition comprising a product produced by mixing in amounts effective to stabilize vinyl halide resins:
- (i) a mono- or diorganotin compound wherein at least one atom bonded to tin is a halogen; and
  - (ii) a mercapto alkanol ester of a monocarboxylic acid.
- 297. The composition of claim 296, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

#### RCOOR'SH

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 298. The composition of claim 297, wherein R contains 6 to 38 carbon atoms.
- 299. The composition of claim 298, wherein R contains 8 to 18 carbon atoms.
- 300. The composition of claim 297, wherein R' contains 2 to 6 carbon atoms.
- 301. The composition of claim 296, wherein the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.
- 302. The composition of claim 296, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl nyristate, mercapto ethyl palmitate and mercapto ethyl myristate.

FINALDAN, HENDERSON
FARASON GARRETT
6 DINNER
1000 I STREET NA
1400 I STREE

303. The composition of claim 296, further comprising a vinyl halide resin.

JEAN-.√ES CHENARD et al. Serial No. 07/570,759

- 304. The composition of claim 303, wherein the product produced by mixing the organotin compound and the mercapto alkanol ester of a monocarboxylic acid are present in an amount effective to stabilize the vinyl halide resin against heat or light.
- 305. The composition of claim 304, wherein the mercapto alkanol ester of a monocarboxylic acid is present in the range of 0.1 % wt. to 5 % wt. of the vinyl halide resin.
- 306. The composition of claim 303, wherein the vinyl halide resin is polyvinyl chloride.
- 307. A method of stabilizing a vinyl halide resin comprising adding to the vinyl halide resin in an amount effective to stabilize the vinyl halide resin a product produced by mixing:
- (i) a mono- or diorganotin compound wherein at least one atom bonded to tin is a halogen; and
  - (ii) a mercapto alkanol ester of a monocarboxylic acid.
- 308. The method of claim 307, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

#### RCOOR'SH

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 309. The method of claim 308, wherein R contains 6 to 38 carbon atoms.
- 310. The method of claim 309, wherein R contains 8 to 18 carbon atoms.
- FINNEDAN HENDERSON FARASON GARRETT 6 DINNER 1200 I STREET NIL TRANSCOND DE 20006 1 202 403 4000
- 311. The method of claim 307, wherein  $R^\prime$  contains 2 to 6 carbon atoms.

- 312. The method of claim 30%, wherein the monocarpoxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.
- 313. The method of claim 30%, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.
- 314. The method of claim 30%, wherein the vinyl halide resin is polyvinyl chloride.
- 315. In a composition for stabilizing a vinyl halide resin containing a mono- or diorganotin compound wherein at least one atom bonded to tin is a halogen, the improvement comprising the addition of a mercapto alkanol ester of a monocarboxylic acid.
- 316. The composition of claim 315, wherein the mercapto alkanol ester of a monocarboxylic acid has the formula:

wherein R is a linear or branched alkyl or alkenyl, aryl or aralkyl containing at least two carbon atoms; and R' designates a  $\rm C_2$  to  $\rm C_{18}$  alkylene.

- 317. The composition of claim 316, wherein R contains 6 to 38 carbon atoms.
- 318. The composition of claim 317, wherein R contains 8 to 18 carbon atoms.
- 319. The composition of claim 316, wherein R' contains 2 to 6 carbon atoms.

ENNEGAN, HENDERSON FARASON, CARRETT 6 DENNER 1222 I STREET & 2 STREETS 22 22223 1224 AUG 4222

- 320. The composition of claim 315, wherein the monocarboxylic acid is selected from the group consisting of caprylic, pelargonic, capric, undecanoic, lauric, myristic, palmitic, stearic, isostearic, and mixtures thereof.
- . 321. The composition of claim 315, wherein the mercapto alkanol ester of a monocarboxylic acid is selected from the group consisting of mercapto ethyl stearate, 3-thio-glyceryl myristate, mercapto ethyl palmitate and mercapto ethyl myristate.
- 322. The composition of claim 315, wherein the mercapto alkanol ester of a monocarboxylic acid is present in the range of 0.1 % wt. to 5 % wt. of the vinyl halide resin to be stabilized.
- 323. The composition of claim 315, wherein the vinyl halide resin is polyvinyl chloride.

FINNELAN, HIS DEMON FAA-4DE CARRETT 5 DINNER 1000 1 STATE N -45015 TO 10 2000 1 201 409 4001